

(19)



JAPANESE PATENT OFFICE

PATENT ABSTRACTS OF JAPAN

(11) Publication number: **03024057 A**

(43) Date of publication of application: **01.02.91**

(51) Int. Cl

**C07D211/40**  
**A61K 31/445**

(21) Application number: **01158162**

(71) Applicant: **TOSOH CORP**

(22) Date of filing: **22.06.89**

(72) Inventor: **NAKANO KOICHI**  
**HASHIMOTO HIRONOBU**

**(54) POLYHYDROXYPIPERIDINES AND PRODUCTION THEREOF**

removed by a catalytic reduction, thus obtaining the objective compound of formula I.

(57) Abstract:



COPYRIGHT: (C)1991,JPO&Japio

NEW MATERIAL: Compounds of formula I (R<sub>1</sub> is H or methyl; One of R<sub>2</sub> and R<sub>3</sub> is H and the other is OH).

EXAMPLE:

2-O-Benzyl-3,4,6-tri-O-acetyl-5-O-trimethylsilyl-D-allono nitrile.

USE: A glycosidase inhibitor.



PREPARATION: A ribofuranoside derivative of formula II [One of R<sub>4</sub> and R<sub>5</sub> is H and the other is alkoxy or formula III (X is R, CH<sub>3</sub>, OCH<sub>3</sub> or Cl); One of R<sub>6</sub> and R<sub>7</sub> is H and the other is acyloxy, etc.; R<sub>8</sub> is acyloxy, etc.; R<sub>9</sub> is acyloxy, azide, etc.] and an arabinofuranoside derivative are reacted with cyanotrimethylsilane in the presence of a Lewis acid and the resultant compound is then subjected to ring opening and carbon increase to obtain a compound of formula IV. The trimethylsilyl group of the resultant compound is substituted for a suitable elimination group and the cyano group thereof is subjected to ring closure by reduction to obtain a compound of formula V. Protective groups of the obtained compound of formula V are